

Birational models for varieties of Poncelet curves

Matei Toma

Universität Osnabrück
Fachbereich Mathematik/Informatik
49069 Osnabrück, Germany
and

Institute of Mathematics of the Roumanian Academy
P.O.Box 1-764, RO-70700 Bucharest, Romania.

We consider Poncelet pairs (S, C) , where S is a smooth conic and C is a degree- c plane curve having the Poncelet property with respect to S . We prove that for $c > 4$ the projection $(S, C) \mapsto C$ is generically one-to-one and use this to describe a birational model of the variety of Poncelet curves for c odd.

Introduction

A pair (S, C) consisting of a smooth conic S and a curve C of degree c in the complex projective plane will be called a Poncelet pair if there exist $c + 1$ tangents to S whose $\binom{c+1}{2}$ intersection points lie on C . C will then be called Poncelet related to S or simply a Poncelet curve. A theorem of Darboux ([3]) states that in this case there is an infinity of such sets of $c + 1$ tangents.

The Poncelet curves reappeared in a natural way in the study of stable vector bundles on projective spaces, e.g. in [1], [5], [2], [11]. Motivated by these developments, Trautmann gave a modern description of their geometry in [12].

In this paper we consider the variety Pon_c of Poncelet curves of degree c . It is the image of the space of Poncelet pairs (S, C) through the second projection. We show that for $c \geq 5$, this projection is birational. This fails for $c \leq 3$, simply because of dimension reasons, and has been proved for $c = 4$ by Le Potier via the study of divisors on the corresponding moduli space of

$rank - 2$ semi-stable sheaves on $\mathbb{P}^2$, ([8]). As a corollary, we get the rationality of Pon_c for every c .

The generic injectivity of the above mentioned projection brings some evidence for (and maybe could help proving) its generalizations to:

- Barth's morphism which associates to a stable $rank - 2$ vector bundle of type $(0, c)$ on $\mathbb{P}^2$ its curve of jumping lines ([1]), and
- the restriction of the previous to Hulsbergen bundles. (The image will be the variety of Darboux curves, cf. [1], [4]).

In the last paragraph we describe a projective birational model of Pon_c when c is odd.

§ 1 Preliminaries

For the proofs of the facts stated in this paragraph as well as for further descriptions we refer the reader to [12].

Let W be a 3-dimensional complex vector space, S a smooth conic in $\mathbb{P}^2 = \mathbb{P}(W)$, c a positive integer and Λ a pencil of degree- $(c + 1)$ divisors on S . We consider the intersection points of the tangents to S at the points of some member D of Λ . When D moves in Λ , these intersection points describe a curve $C = C(\Lambda)$ in $\mathbb{P}^2$. It is clear that the tangents at the base points of Λ will be components of C . We require that their multiplicity in C be equal to that of the base points in D . Then C has degree c , is called **Poncelet related to S** or simply a **Poncelet curve** and (S, C) is called a **Poncelet pair**.

That this definition is equivalent to that of the introduction follows from the

Theorem. (*Darboux*) *Let S be a smooth conic and C a curve of degree c in $\mathbb{P}^2$. If there exist $c + 1$ tangents to S such that their intersection points lie on C , then C is Poncelet related to S .*

The base points of Λ correspond to tangents to S which are components of $C(\Lambda)$ and the residual curve (obtained by eliminating these components) is again Poncelet related to S and associated to the pencil obtained from Λ by subtracting its base points.

For Λ generic $C(\Lambda)$ is smooth.

Let's denote by $G_S(1, c+1)$ the Grassmannian of pencils of degree- $(c+1)$ effective divisors on S . The map

$$\begin{aligned} G_S(1, c+1) &\rightarrow \mathbb{P}(S^c W^*) = \mathbb{P}^{\binom{c+2}{2}-1} \\ \Lambda &\mapsto C(\Lambda) \end{aligned}$$

coincides with the Plücker embedding (for a suitable choice of coordinates, [12]). Thus, there is a 1:1 correspondence between $G_S(1, c+1)$ and the variety $Pon_{c,S}$ of curves C which are Poncelet related to S .

Let U be the open set of $\mathbb{P}(S^2 W^*)$ which parameterizes smooth conics in $\mathbb{P}^2$, $I_U \subset U \times \mathbb{P}^2$ the tautological conic bundle over U , $I_U^{(c+1)}$ the Hilbert scheme of degree- $(c+1)$ effective divisors in the fibers of $I_U \rightarrow U$ (which is a $\mathbb{P}^{c+1}$ -fibration over U) and $G_U(1, c+1)$ the relative Grassmannian of lines in the fibers of $I_U^{(c+1)} \rightarrow U$. We shall identify $G_U(1, c+1)$ with the subvariety $Pon_{c,U}$ of Poncelet pairs in $U \times \mathbb{P}(S^c W^*)$. (The equations of $Pon_{c,U}$ are given in [12]).

We shall examine the second projection $pr_2 : Pon_{c,U} \rightarrow \mathbb{P}(S^c W^*)$. Its image is the variety of degree- c Poncelet curves Pon_c .

We end this paragraph with some elementary remarks on Poncelet curves to be used later:

Remarks. Let S be a smooth conic.

- 1.1 If C is Poncelet related to S , then the only components of C which may be multiple are tangent lines to S . (Apply Bertini's theorem to the associated pencil, Λ .)
- 1.2 If $C_1 + C_2$ and $C_1 + C'_2$ are two elements of $Pon_{c,S}$ such that C_2 and C'_2 have no common component, then C_1 is Poncelet related to S . (Start from a point on C_1 , draw tangents to S and continue this procedure from the intersection points with C_1 . The process stops after a finite number of steps. If there exist intersection points of the drawn tangents not on C_1 , then these points lie on a common component of C_2 and C'_2).
- 1.3 Let $C_1, C_1 + C_2$ be Poncelet related to S with associated pencils Λ_1 and Λ , respectively. If Λ has no base points ,

then the induced morphism $\varphi_\Lambda : S \rightarrow \Lambda^* \cong \mathbb{P}^1$ factorizes through $\varphi_{\Lambda'} : S \rightarrow \Lambda'^*$. In particular, $\deg C_1 + 1$ divides $\deg(C_1 + C_2) + 1$.

- 1.4 If C is a singular conic Poncelet related to S , then one of its components is tangent to S . (Follows from 1.3).

§ 2 Generic injectivity of pr_2

Theorem. *When $c \geq 5$, the projection $Pon_{c,U} \rightarrow Pon_c$ is birational.*

Remarks.

- 2.1 The corresponding statement for $c = 4$ (and more) is proven in [8].

- 2.2 $Pon_{c,U}$ is rational ([6], Prop.2.2), hence so is Pon_c too.

The theorem follows by a dimension count out of the following

Proposition. *Let S, S' be distinct smooth conics. Then*

- (A) *for $c \geq 3$, $\dim(Pon_{c,S} \cap Pon_{c,S'}) \leq c$.*
- (B) *$Pon_{5,S} \cap Pon_{5,S'}$ is at most 4-dimensional at points represented by smooth quintics.*

Indeed, let us estimate the dimension of $\tilde{G}_S := \overline{pr_2^{-1}(Pon_{c,S})} \setminus G_S(1, c+1)$. By part (A) of the Proposition, $\dim \tilde{G}_S \leq c + 5$. Thus, for $c \geq 6$ we have $\dim pr_2(\tilde{G}_S) < 2c = \dim Pon_{c,S}$ proving the theorem in this case.

When $c = 5$, we use part (B) for a slightly modified $\tilde{G}_S$, $\tilde{G}_S := \overline{pr_2^{-1}(Reg_{c,S})} \setminus G_S(1, c+1)$, where $Reg_{c,S}$ is the set of smooth Poncelet curves related to S .

In order to prove the Proposition we look at the linear projection of $Pon_{c,S} \cap Pon_{c,S'}$ with center $L_d := \{\text{degree-}c \text{ curves in } \mathbb{P}(W) \text{ allowing } d \text{ as a component}\} \subset \mathbb{P}(S^c W^*)$, for some common tangent line d to S and S' .

Let d be the tangent at P to the smooth conic S . Then $Pon_{c,S} \cap L_d \cong Pon_{c-1,S}$ corresponds to degree- c pencils on S having P as base point (or equivalently, to the Schubert variety on $G_S(1, c+1)$ of lines contained in a hyperplane).

Let further D be an element of $|\mathcal{O}_S(c+1)|$ containing P and $L_D \subset G_S(1, c+1)$ the Schubert variety of lines through D . Then $L_D \cong \mathbb{P}^c$ is linear in $\mathbb{P}(S^c W^*)$ and cuts on L_d a $(c-1)$ -dimensional linear subspace, (the L_{D-P} of $G_S(1, c)$). Thus $L_D \cup L_d$ spans a c -dimensional linear subspace of $\mathbb{P}(S^c W^*)$.

Conversely, for every c -codimensional linear subspace F which contains L_d , there exists some D as above such that $F \cap Pon_{c,S} = L_D \cup (Pon_{c,S} \cap L_d)$.

For two distinct divisors D, D' as above, L_D and $L_{D'}$ cut themselves in exactly one point lying on L_d . In particular, the projection of center L_d displays $Pon_{c,S} \setminus L_d$ as a vector bundle over $|\mathcal{O}_S(c)|$.

Let now V be an irreducible component of $Pon_{c,S} \cap Pon_{c,S'}$ not contained in L_d .

Let $\pi : V \setminus L_d \rightarrow |\mathcal{O}_S(c)|$ be the restriction to V of the above projection, $t := \dim \pi(V \setminus L_d)$, and f the dimension of a generic fiber of π . Note that the fibers of π are linear.

Lemma 2.1 *When $c \geq 2$, up to finitely many exceptions, the fibers of π are at most $(c-1)$ -dimensional.*

Proof. We choose a general hyperplane in $|\mathcal{O}_S(c)|$ and show that the fibers of π over its points cannot be c -dimensional. Indeed, take P_1 on S such that it doesn't belong to a common tangent to S and S' . The divisors containing P_1 form a hyperplane in $|\mathcal{O}_S(c)|$. Take any such divisor and add P to it in order to obtain a divisor $D = P + P_1 + P_2 + \dots + P_c$ in $|\mathcal{O}_S(c+1)|$. Take further a pencil with base points $P_1, P_2, \dots, P_{c-1}$ and containing D . Its associated Poncelet curve is the union of the tangents at $P_1, \dots, P_{c-1}$ to S and a line through the intersection point of d with the tangent to S at P_c . By Remarks 1.2 and 1.3 not all such curves are Poncelet related to S' . $\square$

Proof of part (A) of the Proposition

We argue by induction on c .

Let $c = 3$ and suppose that $t + f > 3$. Then $t \geq 2$ by Lemma 2.1. We pick then a divisor $D = P + P_1 + P_2 + P_3$ on S and look at the fiber of π contained in L_D .

We get a divisor $D' = P' + P'_1 + P'_2 + P'_3$ on S' such that this fiber is exactly $L_D \cap L_{D'} \setminus L_d$. ($L_{D'}$ is to be considered with respect to S'). Note that P_1, P_2 or P_1, P'_2 may be freely chosen so that this fiber be non empty.

Denote by d_i, d'_i the tangents at P_i to S and at P'_i to S' , respectively, $i \in \{1, 2, 3\}$. We index our $P'_i - s$ so that $d \cap d_i = d \cap d'_i$.

Choose now P_1, P'_2 such that the point $d_1 \cup d'_2$ does not lie on a common tangent to S and S' . In particular, it will not lie on both d_3 and d'_3 , say not on d'_3 .

The Poncelet curves in $L_{D'} \cap L_D$ touch d'_2 at its intersection points with d, d'_1 and d'_3 . (If one of the P'_i had multiplicity $(k+1)$ in D' then the condition on the corresponding Poncelet curves would be to touch S at P'_i with multiplicity k , by [12], Prop. 2.6). If we impose to such a curve that d_1 be a component (which we may do since $f \geq 1$), we get that d'_2 must be a component too. By Remark 1.4 the third component of our cubic is tangent to both S and S' , so $f = 1$ and $t = 3$. This allows a free choice on P'_3 too and we get that the tangents through P'_2 and P'_3 to S' form a conic which is Poncelet related to S . But this contradicts Remark 1.4.

Let now $c > 3$.

Case (A1) $t > \frac{c}{2}, f \geq 1$.

We choose as above $D = P + P_1 + \dots + P_c$ and get in the same way a $D' = P' + P'_1 + \dots + P'_c$ such that $L_D \cap L_{D'} \setminus L_s \neq \emptyset$. In doing this $P_1, \dots, P_t$ are free.

We claim that the choice may be done such that d_1 intersects $d'_2, \dots, d'_t$ away from the vertices of the polygon $d, d'_1, \dots, d'_c$ (with the obvious extension of notation). If this were not true then there would exist a point, say P'_{t+1} , in a “bad position”, e.g. such that $d_1 \cap d'_2 \cap d'_{t+1} \neq \emptyset$. We look then at P_2 instead of P_1 and may find a second point P'_{t+2} , in a bad position, and continue in this way. The assumption $t > \frac{c}{2}$ shows now that our claim holds.

Next, from $f \geq 1$ it follows that we may degenerate a curve

C in our fiber, so that d_1 and hence also $d'_2, \dots, d'_t$ become its components. By Remarks 1.2-4 the other components of C are not all tangent lines to S , otherwise we move the lines d'_i , $2 \leq i \leq t$. In particular $t < c$.

If $f \geq 2$, then we may still move our curve C (in its class), fact which combined with Remark 1.3 shows the existence of a pair of degrees (a, b) such that $t - 1 \leq a < b < c$ and $a + 1$ divides $b + 1$. But this would imply $t \leq \frac{c}{2}$, which proves that $f = 1$ and $f + t \leq c$.

Case (A2) $f > \frac{c}{2}$

The proof is similar to that of Lemma 1.

We may first assume that the general $D = P + P_1 + \dots + P_c$ in the image of π doesn't contain any tangency point of a common tangent to S and S' excepting P . Otherwise we restrict our attention to the subvariety of Poncelet curves containing this tangent and apply the induction hypothesis.

In the fiber of π over D we require that the tangents $d_1, \dots, d_{f-1}$ at $P_1, \dots, P_{f-1}$ to S be components of the Poncelet curves C . Then by Remarks 1.3 and 1.4 there exists a pair of degrees (a, b) with $f - 1 \leq a < b \leq c$ and $a + 1$ divides $b + 1$. This forces $f = \frac{c+1}{2}$ proving part (A) of the Proposition. $\square$

Proof of part (B) of the Proposition

Keeping our previous notations we assume that $t + f = c = 5$.

Case (B1), $t = 5$.

Since in this case π is dominant we find for any choice of points $P_1, \dots, P_c$ on S , an element in $L_D \cap V$, where $D = P + P_1 + \dots + P_c$. Take P_1 on S and P'_1 the corresponding point on S' (draw the tangent d'_1 from $d \cap d_1$ to S'). Choose further a point Q on d'_1 and draw tangents from Q to S . Let these tangents be d_2, d_3 . A curve in $L_D \cap V$ has to contain Q and the intersection points of d'_1 with $d, d'_2, \dots, d'_c$. Thus this curve has d'_1 as a component. In the same way, choosing d_4 through $d'_2 \cap d_1$ fixes d'_2 as a component of C . We may still move our point P_5 , and since $d'_1 \cup d'_2$ is not in $Pon_{2,S}$, we get by Remarks 1.2 and 1.3 degrees a, b with $3 \leq a < b \leq 6$ and $a + 1 \mid b + 1$. This is absurd so case (B1) cannot occur.

Case (B2), $t = 4$.

As in case (A1) we may fix the components $d_1, d'_2, \dots, d'_t$ as we wish. The last component of our Poncelet curve is not tangent to S as already remarked in (A1). Since we may consider d'_2, d'_3 fixed and move d'_4 we get again a contradiction of Remarks 1.2 and 1.4. Case (B2) is thus excluded.

Case (B3), $t = 3$, reduces itself to (A1).

Case (B4) $t = 2$.

We pick as usual two arbitrary tangent lines to S and look at the curves in V containing them. They form an at least 1-dimensional subvariety. Since these two tangents do not form a Poncelet conic with respect to S' we get degrees $3 \leq a < b \leq c$ such that $a + 1$ divides $b + 1$. This implies $c \geq 7$ and excludes this case too.

Case (B5), $t=1$.

If we find some $D = P + P_1 + \dots + P_c$ in the “image” of π containing two points which don't lie on common tangents to S and S' we shall argue as in (B4).

Let's assume then that $P_2, \dots, P_c$ are fixed, all lying on common tangents. We have free choice on P_1 . Imposing that d_1 become a component, we are left with an $(f - 1)$ -dimensional family of curves C in L_{D-P_1} , Poncelet related to S , and such that $C + d_1$ belongs to $Pon_{c,S'}$. Applying to this family the same procedure with a different P_1 , we get a 1-dimensional family of curves C with the above properties with respect to both choices of P_1 . The curves C do not consist only of tangents to S' , since these would have to touch S' only at $P'_2, \dots, P'_c$. We thus get a contradiction of Remarks 1.2-4 when applied to $Pon_{c,S'}$.

Case (B6), $t = 0$.

In this case there exist a divisor $D = P + P_1 + \dots + P_c$ on S such that $V = L_D$. The argument from (B4) shows that at most one point, P_1 say, is not on a common tangent to S and S' . By requiring that $d_3, \dots, d_6$ be components we reduce ourselves to the case $c = 2 = f$, $D = P + P_1 + P_2$. But this is excluded by showing that d_2 has to be a component of C .

If all points of D are tangency points of common tangents to S and S' , then either two of them are multiple in D or one of them has multiplicity bigger than 3 in D . In both cases the associated Poncelet curves are all singular by [12], Proposition 5.1. $\square$

§ 3 A projective birational model of $\overline{Pon}_c$ for c odd

For odd c , $G_U(1, c+1)$ is the relative Grassmannian of 2-dimensional subspaces of a rank- $(c+2)$ vector bundle over U . This vector bundle and thus also $G_U(1, c+1)$ may be extended to the whole of $\mathbb{P}(S^2W^*)$. When $c \geq 5$ one obtains a birational map from this extended relative Grassmannian to $\overline{Pon}_c$.

In this paragraph we describe the linear system which induces this birational map and its base locus as a set. (Note that the knowledge of the full scheme - structure of this base-locus would allow one to compute the degree of $\overline{Pon}_c$). Our method is to consider the stable rank-two vector bundles on $\mathbb{P}(W)$ associated to Poncelet curves and their curves of jumping lines.

Throughout this paragraph c is assumed to be odd and bigger than 4.

The extension of $G_U(1, c+1)$ to $\mathbb{P}(S^2W^*)$ is obvious. Take $I \subset \mathbb{P}(S^2W^*) \times \mathbb{P}(W)$ the incidence variety “points of conics”, $p_1 : \mathbb{P}(S^2W^*) \times \mathbb{P}(W) \rightarrow \mathbb{P}(S^2W^*)$ the first projection, $\mathcal{V} := p_{1,*}\mathcal{O}_I\left(0, \frac{c+1}{2}\right)$ and G the relative Grassmannian of 2-dimensional subspaces in the fibers of $\mathcal{V}$. G is then a compactification of $G_U(1, c+1)$.

Consider now a Poncelet pair (S, C) with S smooth and $C = C(\Lambda)$ with base-point-free pencil $\Lambda \subset \left|\mathcal{O}_S\left(\frac{c+1}{2}\right)\right|$. (In this paragraph we use the notation $\mathcal{O}_S\left(\frac{c+1}{2}\right) := \mathcal{O}_{\mathbb{P}(W)}\left(\frac{c+1}{2}\right)|_S$). Λ induces a surjective morphism

$$\mathcal{O}_{\mathbb{P}(W)}^2 \rightarrow \mathcal{O}_S\left(\frac{c+1}{2}\right).$$

Let F be its kernel. Then F is a stable rank-2 vector bundle on $\mathbb{P}(W)$ with $c_1(F) = -2$, $c_2(F) = c+1$. and its jumping lines are exactly the lines joining the points of some divisor of Λ (see [12], 4.1). Thus, the curve $C' \subset \mathbb{P}(W^*)$ of jumping lines of F is Poncelet related to the conic S' dual to S .

We shall make F fit into a flat family of coherent rank-2 sheaves

over G , and examine the birational map associating the curves of jumping lines to the fibers of this family.

We start with some Lemmata.

Lemma 3.1. *Let $0 \rightarrow E' \rightarrow \mathcal{O}_{\mathbb{P}^2}^N \rightarrow E'' \rightarrow 0$ be an exact sequence with E', E'' coherent torsion-free, indecomposable sheaves on $\mathbb{P}^2$ and $c_1(E') = -1$. Then $N \leq 3$ and E', E'' are slope-stable.*

Proof. E'' torsion-free implies E' locally-free. If E' were not slope-stable, there would exist a locally free subsheaf F of E' with $c_1(F) = 0$. Standard arguments on slope-stable vector bundles (cf. [7], V 8.3) show that F would be a direct summand in E' , contradicting the hypothesis.

Similarly, E'' must be slope-stable.

Let $r' := \text{rank } E', r'' := \text{rank } E$.

Suppose $r' = 1$. Then $E' = \mathcal{O}_{\mathbb{P}^2}(-1)$ and the morphism $\mathcal{O}_{\mathbb{P}^2}(-1) \rightarrow \mathcal{O}_{\mathbb{P}^2}^N$ is induced by N sections in $\mathcal{O}_{\mathbb{P}^2}(1)$. Since $\Gamma(\mathcal{O}_{\mathbb{P}^2}(1))$ is 3-dimensional we may choose when $N > 3$ a basis for $\mathcal{O}_{\mathbb{P}^2}(1)^N$ such that $N - 3$ components of $\mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(1)^N$ vanish. But then E'' would split.

The case $r'' = 1$ is treated in the same way.

We are left with the situation $r' > 1, r'' > 1$. But now the Bogomolov inequality gives

$$c_2(E') \geq \frac{r' - 1}{2r'} \cdot c_1(E')^2 > 0$$

$$c_2(E'') \geq \frac{r'' - 1}{2r''} \cdot c_1(E'')^2 > 0$$

hence $c_2(\mathcal{O}_{\mathbb{P}^2}^N) = -1 + c_2(E') + c_2(E'') > 0$, a contradiction. $\square$

Lemma 3.2. *Let $\mathcal{K}$ be the kernel of the natural surjective morphism*

$$p_1^* \mathcal{V} \rightarrow \mathcal{O}_I \left(0, \frac{c+1}{2} \right)$$

on $\mathbb{P}(S^2 W^) \times \mathbb{P}(W)$. Then $\mathcal{K}$ is locally free and its restrictions to the fibers of p_1 are slope-stable.*

Proof. The restriction to a fiber of p_1 over some point s representing a conic S gives an exact sequence

$$0 \rightarrow \mathcal{K} \big|_{p^{-1}(s)} \rightarrow \Gamma \left(\mathcal{O}_S \left(\frac{c+1}{2} \right) \right) \otimes \mathcal{O}_{\mathbb{P}(W)} \rightarrow \mathcal{O}_S \left(\frac{c+1}{2} \right) \rightarrow 0.$$

It is enough to check that $K := \mathcal{K} \big|_{p^{-1}(s)}$ is locally free and slope stable.

Checking the locally freeness is easy (compare K to its double dual or just use the fact that $\mathcal{K}$ appears from an elementary transformation!).

Suppose that K were not slope-stable.

Since $c_1(K) = -2$, there would exist a destabilizing subsheaf K' of K with $c_1(K') = -1$. By Lemma 3.1. K' would have at most rank 2, hence the rank of K could not exceed 4.

But rank $K = c+2$ and the Lemma follows from our assumption on c . $\square$

Let $\mathcal{S}$ and $\mathcal{Q}$ be the tautological sub- and quotient bundle on the relative Grassmannian G of 2-dimensional subspaces in the fibers of $\mathcal{V}$. We denote by $\mathcal{S}_{\mathbb{P}(W)}$, $\mathcal{Q}_{\mathbb{P}(W)}$, $\mathcal{K}_G$, $\mathcal{V}_{G \times \mathbb{P}(W)}$ the pullbacks to $G \times \mathbb{P}(W)$ of $\mathcal{S}$, $\mathcal{Q}$, $\mathcal{K}$ and $\mathcal{V}$, respectively. (Subscript will be used in the sequel to indicate pullbacks through obvious maps). Denote further by α the composite morphism

$$\mathcal{Q}_{\mathbb{P}(W)}^\vee \rightarrow \mathcal{V}_{G \times \mathbb{P}(W)}^\vee \rightarrow \mathcal{K}_G^\vee$$

on $G \times \mathbb{P}(W)$ and by $\mathcal{E}$ its cokernel.

Remark. One sees immediately that the fiber of $\mathcal{E}$ over a point of G represented by a Poncelet pair $(S, C(\Lambda))$ with S smooth and Λ base-point-free is just the dual of the kernel F of the natural morphism

$$\mathcal{O}_{\mathbb{P}^2(W)}^2 \rightarrow \mathcal{O}_S \left(\frac{c+1}{2} \right).$$

In particular, it is stable and locally free.

We claim that $\mathcal{E}$ is flat over G . This is a consequence of a local flatness criterion ([9]; 2.2.5) and the following

Lemma 3.3 *The restrictions of α to the fibers of $G \times \mathbb{P}(W) \rightarrow G$ are injective and their cokernels are stable as soon as they are torsion-free.*

Proof. Fix a conic S and a c -dimensional subspace A of $\Gamma(\mathcal{O}_S(\frac{c+1}{2}))^* \cong \Gamma(\mathbb{P}(W), K^\vee)$. The slope-stability of K and Lemma 3.1 imply that the natural morphism $s : A \otimes \mathcal{O}_{\mathbb{P}(W)} \rightarrow K^\vee$ is injective.

If Coker s is torsion-free but not stable, it will admit a rank-1 subsheaf E' with torsion-free quotient and $c_1(E') = 1$, $c_2(E') \leq \frac{c-1}{2}$. In particular, $\dim \text{Ext}^1(E', \mathcal{O}_{\mathbb{P}(W)}) \leq \frac{c-1}{2}$. We get a subsheaf K' of $K^\vee$ with $c_1(K') = 1$, $\text{rank}(K') = c + 1$, and which sits in an exact sequence

$$0 \rightarrow A \otimes \mathcal{O}_{\mathbb{P}(W)} \rightarrow K' \rightarrow E' \rightarrow 0.$$

This leads to a contradiction of the stability of K in view of the following simple fact of homological algebra:

Lemma 3.4 *Let A be a finite-dimensional $\mathbb{C}$ -vector space, $\mathcal{C}$ a coherent sheaf on a compact complex manifold X and*

$$0 \rightarrow A^* \otimes \mathcal{O}_X \rightarrow \mathcal{B} \rightarrow \mathcal{C} \rightarrow 0$$

an extension given by an element

$$\eta \in \text{Ext}^1(\mathcal{C}, A^* \otimes \mathcal{O}_X) \cong \text{Hom}(A, \text{Ext}^1(\mathcal{C}, \mathcal{O}_X))$$

such that $A = A' \oplus A''$ and $\eta|_{A''} = 0$. Then

$$\eta|_{A'} \in \text{Hom}(A', \text{Ext}^1(\mathcal{C}, \mathcal{O}_X)) \cong \text{Ext}^1(\mathcal{C}, A'^* \otimes \mathcal{O}_X)$$

gives rise to a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A'^* \otimes \mathcal{O}_X & \longrightarrow & \mathcal{B}' & \longrightarrow & \mathcal{C} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A^* \otimes \mathcal{O}_X & \longrightarrow & \mathcal{B} & \longrightarrow & \mathcal{C} \longrightarrow 0 \end{array}$$

with injective vertical maps. □

Associating to a fiber of $\mathcal{E}$ over G its divisor of jumping lines defines a rational map $G \rightarrow \mathbb{P}(S^c W)$ in the usual way. Indeed, we consider the standard diagram

$$\begin{array}{ccc} \mathbb{F} & \longrightarrow & \mathbb{P}(W^*) \\ \downarrow & & \\ \mathbb{P}(W) & & \end{array}$$

where $\mathbb{F}$ is the incidence variety “points on lines”, then take its product with G

$$\begin{array}{ccc} G \times \mathbb{F} & \xrightarrow{q} & G \times \mathbb{P}(W^*) \\ p \downarrow & & \\ G \times \mathbb{P}(W) & & \end{array}$$

and look at the relative theta characteristic associated to $\mathcal{E}$,

$$\Theta(\mathcal{E}) := R^1 q_* p^*(\mathcal{E}(-2)).$$

One sees immediately that, for the torsion free fibers, $\mathcal{E}_g$ of $\mathcal{E}$, the corresponding fibers of $\Theta(\mathcal{E})$ are the usual theta-characteristics associated to a stable sheaf on $\mathbb{P}^2$ with even first Chern class (cf. [1]). In particular, their supports are the curves of jumping lines of the $\mathcal{E}_g - s$. Let B be the subset of G over which $\mathcal{E}$ has nontrivial torsion in its fibers. Then we get a morphism

$$G \setminus B \rightarrow \mathbb{P}(S^c W),$$

(see [10]).

From the defining sequence of $\mathcal{E}$ one obtains the following resolution of $\Theta(\mathcal{E})$ on $G \times \mathbb{P}(W^*)$

$$0 \rightarrow \mathcal{Q}_{\mathbb{P}(W^*)}^\vee(-1) \xrightarrow{\varphi} (R^1 p_{1*} \mathcal{K}^\vee)_{G \times \mathbb{P}(W^*)} \rightarrow \Theta(\mathcal{E}) \rightarrow 0.$$

In order to obtain the “support” Z of $\Theta(\mathcal{E})$ one takes the determinant of φ and twists it correspondingly to get:

$$0 \rightarrow \mathcal{L}^{-1} \boxtimes \mathcal{O}_{\mathbb{P}(W^*)}(-c) \rightarrow \mathcal{O}_{G \times \mathbb{P}(W^*)} \rightarrow \mathcal{O}_Z \rightarrow 0$$

where

$$\mathcal{L} := \det \mathcal{Q} \otimes \mathcal{O}_{\mathbb{P}(S^2 W^*)} \left(-\frac{(c-1)(c-3)}{8} \right)_G.$$

Notice that Z is not flat over G . In fact, its fibers over B are 2-dimensional. Thus the rational map $G \rightarrow \mathbb{P}(S^c W)$ above is given by a linear subsystem of $|\mathcal{L}|$ and has base locus B . The structural sheaf of this base locus may be recovered by taking

the push-down on G of the previous exact sequence twisted by (-3) . One obtains for it the presentation

$$\mathcal{L}^{-1} \otimes H^0(\mathcal{O}_{\mathbb{P}(W^*)}(c))^* \rightarrow \mathcal{O}_G \rightarrow R^2\pi_*(\mathcal{O}_Z(-3)) \rightarrow 0,$$

where $\pi : G \times \mathbb{P}(W^*) \rightarrow G$ is the projection.

Finally we want to identify B .

Let $\psi : \mathbb{P}(W^*) \times \mathbb{P}(W^*) \rightarrow \mathbb{P}(S^2W^*)$ be the map induced by the product, $p' : \mathbb{P}(W^*) \times \mathbb{P}(W^*) \times \mathbb{P}(W) \rightarrow \mathbb{P}(W^*) \times \mathbb{P}(W^*)$ the projection,

$$I_i = \{(l_1, l_2, x) \in \mathbb{P}(W^*) \times \mathbb{P}(W^*) \times \mathbb{P}(W) \mid x \in l_i\}, \quad i \in \{1, 2\},$$

the incidence varieties and $\mathcal{W} := p'_*(\mathcal{O}_{I_1}(0, -1, \frac{c-1}{2}))$.

Proposition. *$\mathcal{W}$ is a subbundle of $\psi^*\mathcal{V}$ and the natural map from the Grassmannian of 2-dimensional subspaces in the fibers of $\mathcal{W}$ to G is an embedding whose image is B . In particular, B is smooth of dimension $c + 5$.*

Proof. We examine when a fiber of $\mathcal{E}$ over G admits nontrivial torsion. Fix a conic S and a 2-dimensional subspace L of $\Gamma(\mathcal{O}_S(\frac{c+1}{2}))$. If K and E are the fibers of $\mathcal{K}$ and $\mathcal{E}$ over the corresponding point of G we have exact sequences:

$$\begin{aligned} 0 \rightarrow \left(\Gamma\left(\mathcal{O}_S\left(\frac{c+1}{2}\right)\right) / L \right)^* \otimes \mathcal{O}_{\mathbb{P}(W)} &\rightarrow K^\vee \rightarrow E \rightarrow 0, \\ 0 \rightarrow E^\vee \rightarrow L \otimes \mathcal{O}_{\mathbb{P}(W)} &\xrightarrow{\epsilon} \mathcal{O}_S\left(\frac{c+1}{2}\right) \rightarrow \mathcal{E}xt^1(E, \mathcal{O}_{\mathbb{P}(W)}) \rightarrow 0. \end{aligned}$$

where ϵ is the evaluation morphism. Since K is slope-stable, E may only have purely 1-dimensional torsion, and this happens exactly when $\mathcal{E}xt^1(E, \mathcal{O}_{\mathbb{P}(W)})$ has 1-dimensional support or equivalently, when ϵ fails to be “generically surjective on some component of S ”. More precisely, if this is the case, then $S = S_1 + S_2$ and the composite morphism

$$L \otimes \mathcal{O}_{\mathbb{P}(W)}\left(\frac{c+1}{2}\right) \xrightarrow{\epsilon} \mathcal{O}_S\left(\frac{c+1}{2}\right) \rightarrow \mathcal{O}_{S_2}\left(\frac{c+1}{2}\right)$$

vanishes. Then ϵ factors through $\mathcal{O}_{S_1}\left(\frac{c+1}{2}\right)$ and the pullback of L through ψ will be a 2-dimensional subspace in a fiber of $\mathcal{W}$ over $\mathbb{P}(W^*) \times \mathbb{P}(W^*)$. Indeed, pushing the exact sequence

$$0 \rightarrow \mathcal{O}_{I_1}(0, -1, -1) \rightarrow \psi^*\mathcal{O}_I \rightarrow \mathcal{O}_{I_2} \rightarrow 0$$

twisted by $\left(\frac{c+1}{2}\right)$ from $\mathbb{P}(W^*) \times \mathbb{P}(W^*) \times \mathbb{P}(W)$ down to $\mathbb{P}(W^*) \times \mathbb{P}(W^*)$ gives

$$0 \rightarrow \mathcal{W} \rightarrow \psi^* \mathcal{V} \rightarrow p'_*(\mathcal{O}_{I_2} \left(0, 0, \left(\frac{c+1}{2}\right)\right)) \rightarrow 0.$$

The claims of the Proposition are now easy to check; that the considered morphism is an embedding follows e.g. from a computation of its Jacobian matrix.

□

References

- [1] W. Barth: Moduli of vector bundles on the projective plane, Invent. Math. 42, 63-91, (1977).
- [2] W. Böhmer, G. Trautmann: Special instanton bundles and Poncelet curves, Springer LNM 1273, 325-336, Berlin 1987.
- [3] G. Darboux: Géométrie analytique, Paris 1917.
- [4] G. Ellingsrud, S.A. Strømme: Bott's formula and enumerative geometry, to appear in Journal of the AMS.
- [5] R. Hartshorne,: Stable vector bundles of rank 2 on $\mathbb{P}_3$, Math. Ann. 238, 229-280, (1978).
- [6] A. Hirschowitz, M.S. Narasimhan: Fibrés de 't Hooft spéciaux et applications, Proc. Nice Conf. 1981, 143-163, Birkhäuser 1982.
- [7] S. Kobayashi: Differential geometry of complex vector bundles, Iwanami Shoten and Princeton University Press, Princeton 1987.
- [8] J. Le Potier: Fibrés stables sur le plan projectif et quartiques de Lüroth, preprint 1989.
- [9] H. Matsumura: Commutative ring theory, Cambridge University Press, Cambridge 1986.
- [10] M. Maruyama: Moduli of stable sheaves - generalities and the curves of jumping lines of vector bundles on $\mathbb{P}^2$, Advanced Studies in Pure Math. 1, Kinokuniya and North-Holland, 1983.
- [11] M.S. Narasimhan, G. Trautmann: Compactification of $M_{\mathbb{P}_3}(0, 2)$ and Poncelet pairs of conics, Pacific J. 145, 255-365, (1990).
- [12] G. Trautmann: Poncelet curves and associated theta characteristics, Expo. Math. 6, 29-64, (1988).